

ANTIKOAGULATION UND MAMMABIOPSIE

| MEDIKAMENT | STANZBIOPSIE | VAKUUMSAUGBIOPSIE |
|---|--|--|
| Warfarin (Coumandin®) Phenprocoumon (Marcoumar®) Acenocoumarol (Sintrom®) | INR ≤ 4,0 → OK INR > 4,0 → Stopp → Start 6 – 8 Stunden nachher | INR ≤ 2,5 → OK INR > 2,5 → Stopp → Start 6 – 8 Stunden nachher |
| Apixaban (Eliquis®) Dabigatran (Pradaxa®) Edoxaban (Lixiana®) Rivaroxaban (Xarelto®) | OK | → Stopp 48 Stunden vorher → Start 24 Stunden nachher |
| Acetylsalicylsäure (Aspirin®, Thrombo ASS®) Clopidogrel (Plavix®) Ticagrelor (Brilique®) Prasugrel (Efient®) | OK | <i>Bei dualer Therapie</i> → Stopp 5 Tage (Plavix®, Brilique®) bzw. 7 Tage (Efient®) vorher → Start 6 – 8 Stunden nachher |
| LMWH: Enoxaparin (Lovenox®, Inhixa®) Dalteparin (Fragmin®) | OK | <i>Prophylaktische Dosis</i> → Stopp 12 Stunden vorher → Start 6 – 8 Stunden nachher <i>Therapeutische Dosis</i> → Stopp 24 Stunden vorher → Start 24 Stunden nachher |